

# 511-2017-11-10-depression

Rick Gilmore

2017-11-10 14:29:59

# Today's topic(s)

- Depression
- Planning for student-led presentations

# Depression

- Symptoms
  - Unhappy mood, insomnia, lethargy, loss of pleasure, interest, energy
- Agitation
- Lasting for several weeks or more

# Depression


- Experienced by ~7% Americans in any year
- Prevalence (up to ~20% lifetime)
- Females 2-3x males, higher 40+ years of age
- MZ concordance ~60% vs. DZ ~20% suggests genetic component

# Symptoms, (Mahar et al. 2014)

Table 1.

Symptoms of a **depressive episode**, at least five of which must persist for at least two weeks to meet diagnostic criteria, with depressed mood or **anhedonia** requisite (DSM-V; American Psychiatric Association, 2013).

Depressed mood most of the day, nearly every day
Compromised ability to experience pleasure (anhedonia) or interest in activities most of the day, nearly every day
Feelings of worthlessness or unreasonable guilt nearly every day
Sleep disturbance (insomnia or hypersomnia) nearly every day
Fluctuations in weight or appetite changes nearly every day
Psychomotor agitation or retardation nearly every day
Fatigue nearly every day
Diminished ability to think or concentrate nearly every day
Recurrent thoughts of death or suicidal ideation

Table options 

# Neurobiology of Major Depressive Disorder (MDD)


- Reduced sizes of brain regions
- Hypoactivity
- Pharmacological factors
- Synaptic neurotrophic dysfunction

# Neurological factors

- Reduced hippocampal volumes
- [\(Videbech and Ravnkilde 2004b\)](#) meta-analysis

# (Videbech and Ravnkilde 2004b)


## Left Hippocampus





# (Videbech and Ravnkilde 2004a)


## Right Hippocampus



# Hypoactivity in

- Frontal and temporal cortex
- Anterior cingulate
- Insula
- Cerebellum
- [\(Fitzgerald et al. 2008\)](#)

# (Fitzgerald et al. 2008)




[a] patients v. ctrls, [b] patients on SSRIs, [c] patients v. ctrls (happy stim), [d] patients v. controls (sad stim)

# Baseline hyperactivity (Hamilton et al. 2012)




Structure	Direction of Effect	Talairach Coordinates	Cluster Size (mm <sup>3</sup> )	Number
Pulvinar nucleus	Depressed > Comparison	-15, -24, 8	3,054	1
Pulvinar nucleus	Depressed > Comparison	17, -25, 4	2,514	2

# Valence-specific hyperactivity (Hamilton et al. 2012)



Structure	Direction of Effect	Valence Specific Effect?	Talairach Coordinates	Cluster Size (mm <sup>3</sup> )	Number
Amygdala	Depressed > Comparison	Yes	24, -4, -13	318	1
Dorsal anterior cingulate cortex	Depressed > Comparison	Yes	-2, 30, 20	196	2
Insula and superior temporal gyrus	Depressed > Comparison	Yes	-38, -6, -8	834	3
Precentral gyrus	Depressed > Comparison	Yes	-30, -15, 44	621	-
Middle temporal gyrus	Depressed > Comparison	Yes	-39, -64, 17	440	-
Dorsolateral prefrontal cortex	Comparison > Depressed	Yes	30, 13, 47	1,380	4
Dorsolateral prefrontal cortex	Comparison > Depressed	No	-22, 27, 42	949	-
Caudate body	Comparison > Depressed	No	10, 20, 6	382	5


# Increased connectivity between resting state network regions and dorsal PFC (Sheline et al. 2010)





CCN (yellow); precuneus, part of DMN (pink); and affective division of the ACC (turquoise)

# Altered connectivity ([Cheng et al. 2016](#))

- Resting state fMRI (rsfMRI) in 421 patients with major depressive disorder and 488 control subjects.
- Reduced connectivity between orbitofrontal cortex (OFC) and other areas of the brain
- Increased connectivity between lateral PFC and other brain areas





**A****B**

# Pharmacological factors

- Endocrine
  - Thyroid dysfunction ([Medici et al. 2014](#))
  - Altered cortisol reactivity ([Burke et al. 2005](#))

# Pharmacological factors

- Monoamine hypothesis
  - More: euphoria
  - Less: depression
  - Resperine (antagonist for NE & 5-HT) can cause depression
  - Low serotonin (5-HT) metabolite levels in CSF of suicidal depressives ([Samuelsson et al. 2006](#))

# Drug treatments

- Monoamine oxidase (MAO) inhibitors
  - MAO inactivates monoamines in terminal buttons
  - MAO-I's boost monoamine levels
- Tricyclics
  - Inhibit NE, 5-HT reuptake
  - Upregulate monoamine levels, but non-selective = side effects

# Drug treatments

- Selective Serotonin Reuptake Inhibitors (SSRIs)
  - Fluoxetine (Prozac, Paxil, Zoloft)
  - Prolong duration 5-HT in synaptic cleft
  - Also increase brain steroid production
- Serotonin Norepinephrine Reuptake Inhibitors (SNRIs)

# Cymbalta (SNRI)

Cymbalta Commercial (real one)




# How well do drugs work?

- [STAR\\*D trial](#)
- On SSRI for 12-14 weeks. ~1/3 achieved remission; 10-15% showed symptom reduction.
- If SSRI didn't work, could switch drugs. ~25% became symptom free.
- 16% of participants dropped out due to tolerability issues
- Took 6-7 weeks to show response.

# Who benefits from drug therapy?

- Depends on
  - Early life stress
  - Brain (amygdala) response to emotional faces
- [\(Goldstein-Piekarski et al. 2016\)](#)
- Low-stress + low amygdala reactivity -> > responding
- High stress + high amygdala reactivity -> > responding





[\(Goldstein-Piekarski et al. 2016\)](#)

# Problems with monoamine hypothesis

- Too simplistic
- NE, 5-HT interact
- Drugs fast acting (min), but improvement slow (weeks)

**What do drugs do, then?**


# Ketamine again

- Relieves depressive symptoms relatively quickly ([Berman et al. 2000](#)) and ([Zarate et al. 2006](#))
- Boosts synaptic spine formation ([Li et al. 2010](#)) and reverses effects of induced stress


# Pathway of pathology (Duman and Aghajanian 2012)

- Depression ~ chronic stress (Mahar et al. 2014)
- Stress -> chronic HPA axis activity
- Chronic HPA activity -> neuronal atrophy in hipp & PFC
- Stress & cortisol decrease expression of brain-derived neurotrophic factor (BDNF)
- BDNF boosts neurogenesis
- SSRIs act via BDNF, as do NMDA receptor antagonists (e.g., Ketamine)

# (Duman and Voleti 2012)



# (Frohlich and Van Horn 2014)



# Electroconvulsive Therapy (ECT)

- Last line of treatment for drug-resistant depression
- Electric current delivered to the brain causes 30-60s seizure.
- ECT usually done in a hospital's operating or recovery room under general anesthesia.
- Once every 2 - 5 days for a total of 6 - 12 sessions.



# Electroconvulsive Therapy (ECT)

- Remission rates of up to 50.9% ([Dierckx et al. 2012](#))
- Seems to work via
  - Anticonvulsant (block Na<sup>+</sup> channel or enhance GABA function) effects
  - Neurotrophic (stimulates neurogenesis) effects

# Take home messages

- Multi-level, multi-method, multi-variate approaches essential to understanding mental illness
- Developmental processes across the life span
- Networks all the way down...

# References

Berman, R M, A Cappiello, A Anand, D A Oren, G R Heninger, D S Charney, and J H Krystal. 2000. "Antidepressant Effects of Ketamine in Depressed Patients." *Journal of Clinical Psychopharmacology* 47 (4): 351–54.  
<https://www.ncbi.nlm.nih.gov/pubmed/10686270>.

Burke, Heather M, Mary C Davis, Christian Otte, and David C Mohr. 2005. "Depression and Cortisol Responses to Psychological Stress: A Meta-Analysis." *Psychological Science* 30 (9): 846–56.  
doi:[10.1016/j.psyneuen.2005.02.010](https://doi.org/10.1016/j.psyneuen.2005.02.010).

Cheng, Wei, Edmund T. Rolls, Jiang Qiu, Wei Liu, Yanqing Tang, Chu-Chung Huang, XinFa Wang, et al. 2016. "Medial Reward and Lateral Non-Reward Orbitofrontal Cortex Circuits Change in Opposite Directions in Depression." *Journal of Affective Disorders*, October, aww255. doi:[10.1093/brain/aww255](https://doi.org/10.1093/brain/aww255).

Dierckx, Bram, Willemijn T Heijnen, Walter W van den Broek, and Tom K Birkenhäger. 2012. "Efficacy of Electroconvulsive Therapy in Bipolar Versus Unipolar Major Depression: A Meta-Analysis." *Journal of Affective Disorders* 14 (2): 146–50. doi:[10.1111/j.1399-5618.2012.00997.x](https://doi.org/10.1111/j.1399-5618.2012.00997.x).

Duman, Ronald S, and George K Aghajanian. 2012. "Synaptic Dysfunction in Depression: Potential Therapeutic Targets." *Journal of Neuroscience* 33 (6103): 68–72. doi:[10.1126/science.1222939](https://doi.org/10.1126/science.1222939).

Duman, Ronald S, and Bhavya Voleti. 2012. "Signaling Pathways Underlying the Pathophysiology and Treatment of Depression: Novel Mechanisms for Rapid-Acting Agents." *Journal of Clinical Investigation* 122 (1): 47–56.  
doi:[10.1016/j.tins.2011.11.004](https://doi.org/10.1016/j.tins.2011.11.004).

Fitzgerald, Paul B., Angela R. Laird, Jerome Maller, and Zafiris J. Daskalakis. 2008. "A Meta-Analytic Study of Changes in Brain Activation in Depression." *Journal of Affective Disorders* 29 (6): 683–95.  
doi:[10.1002/hbm.20426](https://doi.org/10.1002/hbm.20426).

Frohlich, Joel, and John D Van Horn. 2014. "Reviewing the Ketamine Model for Schizophrenia." *Journal of Clinical Psychopharmacology* 28 (4): 287–302. doi:[10.1177/0269881113512909](https://doi.org/10.1177/0269881113512909).